



OPEN ACCESS

The role of IL-17A in axial spondyloarthritis and psoriatic arthritis: recent advances and controversies

Dennis G McGonagle,^{1,2} Iain B McInnes,³ Bruce W Kirkham,⁴ Jonathan Sherlock,^{5,6} Robert Moots^{7,8}

Handling editor Josef S Smolen

¹Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, UK

²NIHR Leeds Musculoskeletal Biomedical Research Unit, Chapel Allerton, Leeds Teaching Hospital Trust, Leeds, UK

³Institute of Infection Immunity and Inflammation, University of Glasgow, Glasgow, UK

⁴Rheumatology Department, Guy's and Saint Thomas' NHS Foundation Trust, London, UK

⁵Kennedy Institute of Rheumatology, University of Oxford, Oxford, UK

⁶Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Oxford, UK

⁷Institute of Ageing and Chronic Disease, University of Liverpool, Liverpool, UK

⁸Department of Academic Rheumatology, Aintree University Hospital, Liverpool, UK

Correspondence to

Dr Robert Moots, University of Liverpool Institute of Ageing and Chronic Disease, Liverpool, L9 7AL, UK; r.j.moots@liverpool.ac.uk

Received 8 March 2019

Revised 17 May 2019

Accepted 14 June 2019

Published Online First

5 July 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: McGonagle DG, McInnes IB, Kirkham BW, et al. *Ann Rheum Dis* 2019;**78**:1167–1178.

ABSTRACT

Although the pathogenic mechanisms underlying axial spondyloarthritis (axSpA) and psoriatic arthritis (PsA) are not fully elucidated, several lines of evidence suggest that immune responses mediated by interleukin 17A (IL-17A) play a pivotal role in both diseases. This is best highlighted by the significant clinical efficacy shown with inhibitors of IL-17A in treating axSpA and PsA. Nevertheless, a number of knowledge gaps exist regarding the role of IL-17A in the pathophysiology of spondyloarthritis in man, including its cellular origin, its precise role in discrete disease processes such as enthesitis, bone erosion, and bone formation, and the reasons for the discrepant responses to IL-17A inhibition observed in certain other spondyloarthritis manifestations. In this review, we focus on the latest data from studies investigating the role of IL-17A in ankylosing spondylitis (AS) and PsA that build on existing and emerging scientific knowledge in the field. Key remaining research questions are also highlighted to guide future research.

INTRODUCTION

The spondyloarthritides (SpA) comprise related but phenotypically distinct inflammatory diseases including psoriatic arthritis (PsA), non-radiographic axial spondyloarthritis (nr-axSpA) and radiographic axSpA (ankylosing spondylitis (AS)), arthritis associated with inflammatory bowel disease (IBD), reactive arthritis, juvenile idiopathic arthritis and acute anterior uveitis.^{1–3} The SpA diseases share common immunological and inflammatory components and present with overlapping clinical phenotypes.^{4–7} Indeed, multiple genetic polymorphisms within the interleukin (IL)-23/17 axis have been implicated across SpA.^{8–11} Intriguingly, despite the clinical and genetic similarities, these disorders are showing emergent and unexpected heterogeneity with respect to IL-23/17 axis therapeutic manipulation, a topic addressed later in this article.

IL-17A, a member of the IL-17 superfamily of cytokines, is known to play an important role in SpA manifestations related to the skin, joints and entheses, as reflected by the suppression of disease activity seen with IL-17A inhibitors in psoriasis, PsA and AS.^{12–19} However, in other settings where IL-17 family members have been found at sites of disease, such as gut inflammation and uveitis, IL-17A inhibition is not beneficial.^{20–22} These discrepant responses illustrate the need for clearer understanding of the aetiology of these inflammatory diseases, particularly the role of the IL-17 family in the context of the tissue(s) affected. IL-17 research has accelerated rapidly, with nearly 10 000 articles published on this topic in the last 5 years alone. In

this article we highlight the latest breakthroughs that expand understanding of the role of IL-17A in both homeostasis and in disease in axSpA and PsA.

IL-17A PRODUCTION AND SIGNALLING

The IL-17 superfamily consists of six ligands (IL-17A to IL-17F), which can bind to five receptor subtypes (IL-17RA to IL-17RE). The basic biology of most of the IL-17 superfamily has been reviewed extensively elsewhere.^{23–24} IL-17A, the prototypical ligand, is by far the best characterised member of the IL-17 family and can exist as a homodimer or in a heterodimer with IL-17F and signals through an obligate dimeric IL-17RA and IL-17RC receptor complex.²⁴ On binding to a receptor, IL-17A upregulates inflammatory gene expression either by inducing de novo gene transcription or by stabilising mRNA of pro-inflammatory cytokines and chemokines.²⁴

WELL-DEFINED ROLE OF IL-17A IN HOST DEFENCE

In healthy individuals, IL-17A, as well as other members of the IL-17 family, functions in host defence against a range of bacterial and fungal pathogens at epithelial and mucosal barriers in the skin, colon and airways.^{25–27} Although the exact interplay between the various IL-17 family members is poorly understood, epithelial cell-derived (especially IL-17C) and haematopoietic cell-derived IL-17s (IL-17A and F) may have complementary functions in response to pathogens, with the former predominantly enhancing barrier function and the latter propagating the inflammatory response.²⁸ The IL-23/17 axis co-ordinates barrier function in the skin and the gut, both of which are sites of either physical or chemical stress and are also sites of complex microbial interactions. What might the common denominator be between the IL-23/17 axis and inflammation at the sterile skeletal locations afflicted by SpA-associated pathology? We agree with the assertion that the IL-23/17 axis might be adapted to facilitate homeostasis at these highly mechanically stressed enthesal sites that are prone to microinjury.²⁹

An array of genetic defects in the IL-17 pathway, identified through human translational immunology, collectively point towards a role in antifungal immunity (table 1). Chronic mucocutaneous candidiasis (CMC) is a hallmark of individuals with genetic defects affecting IL-17 immunity, manifesting as recurrent or persistent infections of the skin, nails and mucosae with *Candida albicans*, with or without other clinical signs.³⁰ As can be

Table 1 Diseases characterised by mutations and deficiencies in the IL-23/IL-17 pathway

Disease	Genes/proteins involved	Common infections	Effect on IL-23/IL-17 signalling	References
MSMD	IL-12Rβ1	Mycobacterium tuberculosis, salmonella, candida albicans	Impaired IFN-γ-mediated immunity Decreased IL-17A-producing T cells Impaired IL-12 signalling	176 181 183–187 189
	IL-12β	Mycobacterium tuberculosis, salmonella		
	IFN-γR1	Mycobacterium tuberculosis, salmonella, candida albicans		
	STAT1	Mycobacterium, candida albicans		
	IL-12/23 p40	Mycobacterium tuberculosis, salmonella		
Salmonella infection	IL-23R IL-12/23p40 IL-12Rβ1	Salmonella	Th17 depletion, reduced production of IL-17A	188 190
CMCD	STAT1	Candida albicans, mycobacterium	Reduced production of IL-17A, IL-17F, IL-22	182 183 191 193 202
	CARD9	Candida albicans	No response to IL-17A; reduced response to IL-17F; IL-17E response maintained	
	IL-17RA	Candida albicans Staphylococcus	Impaired neutrophil function	
	IL-17RC	Candida albicans		
	IL-17F	Candida albicans		
	ACT1	Candida albicans Staphylococcus		
APECED	AIRE	Candida albicans	Increased autoantibodies to IL-17A, IL-17F, and IL-22	194 197
HIES	STAT3	Candida albicans, staphylococcus, aspergillus	Increased serum IgE Eosinophilia	49 198 201 203
	DOCK8	Candida albicans, staphylococcus, aspergillus	Impaired development of Th17 cells Reduced IL-17A production	
	TYK2	Staphylococcus, mycobacterium, salmonella		

The genes and proteins involved and the resultant effects on IL-17 signalling that lead to increased susceptibility to certain infections are listed.^{49 176–201} IL-12 and IL-23 share cytokine and receptor subunits and the association with mycobacterial disease is thought to represent an effect on dysregulated IFN gamma production in the IL-12 pathway. Therefore IL-17 blockers or IL-23 p19 subunit blockers are not expected to have a link with mendelian susceptibility to mycobacterial disease.

AIRE, autoimmune regulator; APECED, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy; CARD9, caspase recruitment domain-containing protein 9; CMCD, chronic mucocutaneous candidiasis disease; DOCK8, dedicator of cytokinesis 8; HIES, hyper IgE syndrome; IFN, interferon; IFN-γR1, interferon gamma receptor 1; IL-12Rβ1, interleukin 12 receptor β1; MSMD, Mendelian susceptibility to mycobacterial disease; STAT, signal transducer and activator of transcription; TYK2, tyrosine kinase 2; Th-17, T helper 17 cell.

seen from [table 1](#), these syndromes result from genetic defects affecting several immune processes, with the commonality being that the defects involve more than one cytokine or immune function. The genetic defects that are often shared between the IL-12 and the IL-23 pathway that are upstream of the IL-17 pathway are not linked to fungal infection but may be linked to mycobacterial infections consequent to impaired interferon gamma signalling.³¹

Preclinical and ex vivo studies also implicate IL-17A in immunity against a range of other pathogens including bacteria such as *Escherichia coli*, fungi such as *Cryptococcus neoformans*, parasites such as *Trypanosoma cruzi*, and viruses such as influenza (reviewed in Matsuzaki and Umemura).²⁸ Although as with all drugs that modulate immune response there is the potential for an increased infection risk with IL-17A inhibitors, clinical data show no risk from specific pathogens, with the exception of candidiasis.^{18 19 32–35} Reassuringly, no association between mycobacterial disease and blockade of IL-17A biological activity has been observed in man.³⁴

ROLE OF IL-17A IN SPA

Although IL-17A cytokine expression has been detected in a multitude of autoimmune and autoinflammatory diseases, a key role in psoriasis, PsA and axSpA is evident.

Genetics

Although a detailed examination of the genetic basis of SpA is outside the scope of this article (reviewed in detail in Taams *et al* and Brown *et al*),^{11 36} the strongest association with genetic susceptibility to axSpA and PsA lies within the MHC class I region and in particular the HLA-B27 region.^{8–10} Multiple immunological functions can be altered by these genetic associations, including several relevant to IL-17A signalling through activation of CD8+T cells and CD4+T cells.^{11 37–40} Several single nucleotide polymorphisms in genes directly involved in IL-17 signalling have also been linked to AS and PsA ([figure 1](#)), including variants in the IL-12 p40 subunit, the IL-23 p19 subunit, the IL-23 receptor, IL-17A and IL-17RA.^{41–48} Additional susceptibility variants have been identified in genes encoding IL-17-related signalling molecules including TYK2, TRAF3IP2 and STAT3.^{43 44 47 49–53}

IL-17A production

There has been significant interest in identifying the sources of IL-17A in SpA and a number of innate and adaptive immune system cell types have been implicated (reviewed in Taams *et al*) ([figure 2](#)).¹¹ Increased levels of type 3 innate lymphoid cells (ILC3) have been identified in blood and synovia of patients




Figure 1 Single nucleotide polymorphisms identified in the IL-17 signalling pathway that have been linked to axial spondyloarthritis and psoriatic arthritis. ^aSignificant association shown in European but not Asian populations²⁰⁴; ^bNo risk associated with this SNP shown in certain studies^{204–206}; ^cNo risk associated with this SNP shown in certain studies²⁰⁷; ^dSNP can be associated with risk or protection depending on the specific mutation. AS, ankylosing spondylitis; IL-1R2, interleukin 1 receptor; IL-6R, interleukin 6 receptor; IL17R, interleukin receptors; PsA, psoriatic arthritis; SNP, single nucleotide polymorphism.

with SpA, and these levels correlate with PsA disease activity.^{54–55} Recently, resident populations of both ILC3 and $\gamma\delta$ T cells have been identified at the human enthesis for the first time where they may produce the IL-17A that drives enthesitis, a key early pathological lesion in SpA.^{56–59} Recent studies have also reported

increased numbers of IL-17+invariant natural killer T (iNKT) cells and $\gamma\delta$ T cells in SpA patient peripheral blood.⁶⁰ Indeed, these ROR γ t+innate like T cells, and not conventional T cells, represented about half of all IL-17 producing blood circulating T cells and were further skewed towards IL-17 expressing subsets




Figure 2 Key sources of IL-17A in spondyloarthritis. AS, ankylosing spondylitis; ILC3, type 3 innate lymphoid cell; iNKT, innate natural killer T cell; MAIT, mucosal-associated invariant T cell; PsA, psoriatic arthritis; SpA, spondyloarthritis; Th17, T helper 17 cell; T_{RM}, resident memory T cell.

in synovial fluid samples, as determined by both advanced cytometric methodologies and intracellular cytokine IL-17 staining.⁶⁰ In vivo evidence for enrichment of pathogenic subsets in the joints was recently shown in mannan-induced arthritis in SKG mice, an IL-23/17 axis dependent disease.⁶¹

Tissue-resident memory T cells (T_{RM}) represent approximately 50%–70% of the pool of resident T cells in healthy skin, and can produce a variety of cytokines, including IL-17A.^{62–64} In patients with psoriasis, IL-17-producing CD8+ T_{RM} cells have been identified in non-involved skin and may be involved in recurrence of psoriasis at sites of prior resolution.^{62–64} Efforts are ongoing to investigate the role of T_{RM} cells in tissues affected by SpA. A study in patients with PsA has shown the presence of IL-17 expressing CD4- (CD8+) T cells in the synovium.⁶⁵ However, no studies have been reported in axial disease to date.

Adaptive immune cells are key drivers of chronicity in SpA and as such, are also a major source of IL-17A. The presence of T helper 17 (Th17) cells in SpA is relatively well established.⁶⁶ Increased levels of both Th17 cells and IL-17A are found in skin lesions and the blood of patients with psoriasis as well as the blood and synovial fluid of patients with AS and PsA.^{67–75} IL-17A initiates several feedback-loop mechanisms in SpA leading to increased expansion of Th17 cells and thereby further production of IL-17A.⁷⁶ Evidence suggests that there are distinct subtypes of Th17 cells whose differentiation is dependent on specific combinations of cytokines.⁷⁷ Furthermore, there is likely to be considerable plasticity between Th17 cells and FOXP3+ regulatory T cells.⁷⁸ Identification of the subtype(s) of Th17 cells and their regulation and relevance to axSpA and PsA is an important topic of ongoing research.

Although not found in rheumatoid arthritis synovial fluid, IL-17A-producing conventional CD8+T cells are present in synovial fluid of inflamed joints in patients with AS and PsA where their levels correlate with disease activity.^{71 72 79} IL-17-producing mucosal-associated invariant T (MAIT) cells have been identified in skin and blood from patients with psoriasis⁸⁰ and are also increased in the synovial fluid and blood of patients with AS, where they produce IL-17A in an IL-7-dependent fashion.^{81 82}

It has been suggested that neutrophils contribute to the amplification of the inflammatory response in SpA by producing further IL-17A and although IL-17A-positive neutrophils have been reported in psoriatic skin, the synovium of patients with PsA, and in AS facet joints,^{83–88} the emerging consensus is that neutrophils do not produce IL-17A mRNA or protein, even after strong stimulation with various cytokine combinations.⁸⁹ Similarly, although IL-17A-positive mast cells have been found in synovial tissue from patients with SpA,⁹⁰ the concept is of exogenous IL-17A capture and release, as opposed to synthesis.⁹¹ A recent study indicated that levels of IL-17A were higher in joint-resident mast cells following IL-17A inhibition, which supports the concept of storage of this cytokine under normal tissue homeostasis and mast cell release during inflammation.⁹²

A key research question for the future will be identifying all IL-17A-producing populations, especially at the entheses in SpA. It is noteworthy that measurement of serum levels of IL-17A is likely to be of minimal relevance due to the local tissue responses seen in SpA via IL-17A-producing resident cells.

Enthesitis

Considerable recent developments have occurred in experimental enthesitis research where high mechanical stressing at entheses is associated with local immune system activation.^{59 93 94} Non-SpA-related enthesitis can result from repeated mechanical

strain in healthy individuals (eg, tennis elbow) and usually resolves spontaneously, whereas inflammation in SpA shows chronicity.⁵⁹ The underlying mechanisms behind this pathologically exaggerated immune response, which is driven by a combination of genetic factors and disturbed epithelial barrier function, are starting to be unravelled.^{59 95–97}

Enthesitis is triggered predominantly by an innate immune response. Prostaglandin E2 (PGE2) and IL-23 may be important early mediators, activating resident immune cells to produce IL-17A and other inflammatory cytokines. Indeed, peri-enthesal bone involvement and the often excellent responses observed with NSAIDs incriminate PGE2 in axial disease.⁵⁹ In mice, hepatic expression of IL-23 induces spondyloarthropathy by acting on ROR- γ t+CD3+CD4-CD8- enthesal resident T cells to produce inflammatory mediators including IL-17A.⁹⁸ γ/δ T cells have been shown to constitute the large majority of murine IL-17A producing cells, proliferating at the site of injury, and enhancing bone regeneration.^{99 100} However, although enthesitis appears to be a cardinal lesion in several IL-23/17 axis murine models of inflammatory arthritis,^{101–107} other models have indicated that disease can arise in a T-cell independent manner including that mediated by TNF production from enthesal myeloid and stromal cells.^{94 108–110}

In humans, IL-17A-producing entheses-resident ILC3 and γ/δ T cells have recently been described.^{56–58} Resident myeloid cells that can locally produce IL-23 have also been described,¹¹¹ and their numbers in man may be linked to mechanical load.¹⁰⁸ IL-17A likely acts as an amplifier of enthesitis, inducing several other cytokines by resident mesenchymal cells.^{98 112 113} Prolonged enthesal inflammation leads to new bone formation and also, to a much lesser extent, bone erosion, and is subject to considerable research interest.⁵⁹

Bone damage

Preclinical and clinical data suggest that bone erosion and new bone formation in SpA may occur simultaneously at different anatomical sites,^{114–117} with IL-17A playing a complex role in these processes.

Bone erosion

Numerous preclinical studies have indicated that IL-17A promotes bone resorption in experimental arthritis.^{118–121} Recent efforts have focused on elucidating the mechanisms behind these effects and indicate that IL-17A stimulates receptor activator of nuclear factor- κ B ligand (RANKL) expression and inhibition of Wnt signalling, thereby inhibiting osteoblast activity (figure 3).^{122–128}

Clinical data in patients with PsA show a significant reduction in joint radiographic progression with IL-17A inhibitors versus placebo in the short-term and low long-term rates of radiographic progression.^{13 14 129} Furthermore, recent data from the PSARTROS study showed no progression of catabolic and anabolic bone changes in the joints of patients with PsA treated with secukinumab for 24 weeks.¹³⁰

New bone formation

The precise role of IL-17A in new bone formation in axial SpA and PsA is currently unknown, with contradictory experimental findings observed. Studies favouring a role in new bone formation include data from both animal models and human primary cells (summarised in figure 3). IL-17A has been reported to boost osteogenesis via enhancing osteoblast differentiation from local mesenchymal stem cell populations,^{100 131–133} and the subsequent




Figure 3 The role of IL-17A in bone erosion and bone formation in spondyloarthritis. Adapted from Schett *et al* and Gravalles *et al* [59 223 231]. BMP, bone morphogenetic protein; ILC3, type 3 innate lymphoid cell; RANKL, receptor activator of nuclear factor kappa-B ligand; Th17, T helper 17 cell.

activation of the osteoblasts via activation of the JAK2/STAT3 signalling pathway, which is associated with osteogenesis.¹³¹ IL-17A knockout models have been associated with impaired bone regeneration at both 14 and 21 days post a drill-hole fracture in the femur when compared with wild type mice.¹⁰⁰ Furthermore, in the mycobacterium tuberculosis-induced diseased HLA-B27 transgenic rat model of SpA, IL-17A blockade significantly suppressed pathological new bone formation.¹³⁴ In humans, IL-17A levels are elevated in the days following fracture, which in turn is associated with callus formation.¹³⁵

In contradistinction, cutaneous-restricted overexpression of IL-17A was associated with bone loss in murine models.¹³⁶ Moreover, rat calvarial defects show impaired healing when exposed to IL-17A, combined with significant impairments in osteogenesis in the isolated cells when exposed to IL-17A.¹³⁷ In vivo, IL-17A is associated with osteoclastogenic activation and systemic bone loss in rheumatoid arthritis.^{122 124} Thus, determining the role of IL-17A in new bone formation remains an important avenue of future research.

Pain

The immune system plays a critical role in modulating acute and chronic pain in both the peripheral and central nervous systems.^{138 139} Although pain in SpA is often assumed to be a surrogate marker for inflammation, evidence is emerging to suggest a more complex picture. In axSpA, pain does not always correlate with inflammation or radiographic measures of disease.¹⁴⁰ Furthermore, neuropathic pain as well as inflammatory pain has been observed in patients with AS and PsA.^{141 142}

IL-17A can modulate inflammatory pain by directly increasing nociceptor excitability and potentiating hyperalgesia through the induction of secondary factors.^{139 143–146} Both IL-17RA and IL-17RC are expressed in murine neuronal tissue where they contribute to inflammatory responses.^{147 148} Preclinical studies also suggests a role for IL-17A in neuropathic pain.^{149–152} Clinical data with inhibitors of IL-17A in AS and PsA show rapid and

significant pain reduction,^{153 154} but work to assess whether this represents a reduction in neuropathic as well as inflammatory pain is needed.

Gut inflammation in SpA

The role of IL-17A in IBD and its potential link to the pathogenesis of axSpA and PsA has been the subject of some controversy. Historically, preclinical data investigating the outcome of IL-17A inhibition in mouse IBD models have been inconsistent, with some studies showing disease protection and others showing exacerbation.^{155 156} Clinically, IL-17A inhibition was ineffective in moderate-to-severe Crohn's disease.²⁰ Long-term clinical trial and postmarketing safety data in psoriasis, PsA and AS indicate that the overall incidence of IBD is low, within the expected range in these disorders, and not exacerbated by secukinumab treatment.¹⁵⁷ This highlights one of the pitfalls of translating preclinical data to a clinical setting and has led researchers to reconsider the preclinical IBD models. Nevertheless, long-term data with IL-17A inhibitors in clinical practice are required to investigate this issue further.

The $\gamma\delta$ T cell was the principal source of gut-derived IL-17A in a mouse model of colitis, where IL-17A-dependent regulation of the tight junction protein occludin during epithelial injury was shown to maintain barrier integrity.¹⁵⁸ Mucosal tissues have also emerged as a key physiological site for the differentiation and regulation of Th17 cells.^{77 159 160} A role for ILC3 and innate-like T cells such as iNKT cells and MAIT cells in IBD is also postulated based on their high representation at barrier sites.^{161–163} Putative links have also been suggested between gut inflammation, migration and accumulation of IL-17A-producing ILC3 cells in the joints of patients with AS.¹⁶⁴ A recent study also found that pathogenic bacteria can induce intestinal barrier defects and translocate to systemic organs, triggering autoimmune disease.¹⁶⁵

Uveitis

Like IBD, anterior uveitis in SpA shares common genetic risk factors and the involvement of certain pro-inflammatory

	Secukinumab ^{12–16,20,21,168,232–235}	Ixekizumab ^{17–19,169,236,237}
Approved indications	Psoriasis, PsA, AS	Psoriasis, PsA
MoA	Fully-human anti-IL-17A antibody	Humanised anti-IL-17A antibody
PsA		
Peripheral arthritis	●	●
Enthesitis	●	●
Dactylitis	●	●
Skin	●	●
Nail	●	●
Axial disease	●	▨
Structural progression	●	●
AS		
Signs and symptoms	●	●
Disease activity	●	●
Function	●	●
Structural progression	●	●
Non-radiographic axial SpA	●	●
Psoriasis	●	●
Crohn's disease	●	▨
Uveitis ^a	●	▨

● Effective ● Under investigation ● Not effective ▨ Not assessed

Figure 4 Summary of clinical efficacy with IL-17A inhibitors in spondyloarthritis. ^aNo efficacy shown with secukinumab in non-infectious uveitis; not investigated in anterior uveitis, the form of the disease most common in patients with spondyloarthritis. AS, ankylosing spondylitis; PsA, psoriatic arthritis; SpA, spondyloarthritis.

cytokines. Clinical trials have demonstrated the efficacy of anti-TNF monoclonal antibody therapy in panuveitis or posterior uveitis but clinical trials with inhibitors of IL-17A have failed to meet their primary endpoints for these forms of the disease that are pathophysiologically distinct from anterior uveitis.²¹ Both IL-17A and IL-17F have been detected in anterior uveitis (reviewed in Weinstein and Pepple), but whether they play a critical role is unclear.¹⁶⁶ In secukinumab-treated AS patients there was no evidence suggesting uveitis flares in patients with previous anterior uveitis.¹⁶⁷ Further research is required to extend our understanding of the precise role of IL-17A in the pathogenesis of anterior uveitis.

TARGETING IL-17A IN SPA


The key role played by IL-17A in the pathogenesis of AS and PsA is highlighted by the efficacy shown by inhibitors of IL-17A in clinical trials. Secukinumab, a fully human anti-IL-17A monoclonal antibody, is approved for the treatment of psoriasis, PsA and AS based on the results of several large randomised controlled trials.^{12–16 168} Ixekizumab, a humanised anti-IL-17A antibody, is approved for the treatment of psoriasis and PsA and has shown significant efficacy in two large phase III trials in AS.^{17–19 169 170} The efficacy of IL-17A inhibitors across all manifestations of disease in AS and PsA, including skin, nails, peripheral arthritis, axial disease, dactylitis and enthesitis, highlights the utility of drugs targeting this pathway (figure 4). Inhibitors of IL-17A have also been shown to have an overall favourable long-term safety profile in clinical trials,^{32 33 35} including low rates of serious infections, *Candida* infections

and malignancy, with no evidence of increased suicidality or IBD exacerbation above expected background levels.¹⁵⁷ Nevertheless, the long-term safety of IL-17A inhibitors will need to be monitored in a real-world setting.

WHAT IS THE BASIS FOR DIVERGENT IL-17A AND IL-23 RESPONSES IN AXIAL DISEASE?

IL-23 plays a key role in amplifying and maintaining IL-17A production in many cells, so it was expected that IL-23 inhibitor therapy would have similar results to IL-17A inhibition in axSpA. Interestingly, clinical studies with ustekinumab, an IL-12/-23 inhibitor, in axial SpA were terminated due to lack of efficacy¹⁷¹ and the IL-23 p19 inhibitor risankizumab also failed to show efficacy in AS in a phase II proof of concept study.¹⁷² Conversely, the efficacy of IL-17A inhibition in AS suggests that IL-17A and not IL-23 is the major cytokine mediating disease pathogenesis in axSpA and in this context, IL-17A is likely to be produced in a largely IL-23-independent manner. Understanding the reasons for these divergent roles of IL-23 and IL-17A in the pathophysiology of axSpA is one of the hottest topics in current IL-17A research. Emerging evidence suggests there may be anatomical and immunological differences between axial and peripheral enthesitis and subsequent downstream disease manifestations (figure 5). For instance, there is generally more enthesal soft tissue inflammation or synovio-enthesal complex disease in peripheral enthesitis in PsA,¹⁷³ and more peri-enthesal osteitis in the spine in AS, with this bone proclivity being linked to carriage

A. Anatomic differences between spinal and peripheral entheses



B. Differential cytokine effects in AS, PsA and PsO

	AS	PsA	PsO
IL-23	–	+	++
IL-17A	++	++	++
TNF	+	+	+

C. IL-17A production in spinal entheses




Figure 5 Emergent scheme to explain IL-23/–17 axis pathway divergence in PsA and AS. IL-23 pathway blockade is highly effective in psoriasis but not in AS, which is unexpected given the IL-23 SNPs and related gene SNPs associated with AS. Anatomical differences between entheses in the spine versus peripheral joints could play a role (A). The peripheral skeleton has numerous synovio-entheseal complexes,¹⁷³ which contain abundant myeloid cells, while these cells are rare in the spine. Spinal enthesitis is also associated with peri-entheseal bone disease and osteitis.^{59 173 238} The role of inflammatory cytokines, namely IL-23, IL-17A and TNF α , also differs across the spondyloarthritis (B).^{12 14–16 167 168 172 239–248} IL-17A can be produced by several different sources in spinal entheses (C).^{56–59 74 75 81 82 99 100 158 175 249} Emerging evidence supports the cellular basis for IL-17 production that is independent of IL-23.^{56 57 158 175} Animal models also show that IL-23 has a redundant role once adaptive immunity is primed.¹⁷⁵ Where ++, strong involvement; +, involvement; –, no involvement. AS, ankylosing spondylitis; $\gamma\delta$ T, gamma delta T cells; HLA-B27, human leucocyte antigen B27; IL-17A, interleukin 17A; IL-23, interleukin 23; ILC3, Type three innate lymphoid cells; iNKT, innate natural killer T cell; MAIT, mucosal associated invariant T cell; MSCs, mesenchymal stem cells; PsA, psoriatic arthritis; PsO, psoriasis; Tc17, CD8+T cells; Th17, T helper 17 cells; TNF, tumour necrosis factor α .

of the HLA-B27 gene for axial disease.¹⁷⁴ In terms of control of IL-17 production, IL-23 receptor positive and negative subpopulations of $\gamma\delta$ T cells have been identified in human spinous processes entheses, pointing to a role for IL-23-independent IL-17A production,⁵⁷ and entheses-resident myeloid cells are capable of IL-23 production locally.¹¹¹ Further research is required to investigate the drivers of this process in the future although data in mice indicate that the initiation, but not the persistence, of experimental SpA is dependent on IL-23.¹⁷⁵

CONCLUSION

The IL-17A inhibitors show efficacy in treating multiple facets of SpA, including psoriasis, enthesitis, synovitis, bone erosion, new bone formation and pain, which illustrates the importance of IL-17A in disease pathophysiology. Future research will investigate key remaining gaps, such as the role of human entheses-resident innate and adaptive T cells in SpA and our understanding of IL-23-independent IL-17A production. The ongoing assessment of IL-17A inhibitors in a real-world setting will also be important as these agents become more widely prescribed in clinical practice. Ongoing research efforts will attempt to answer these and other open questions and shed further light on the role of IL-17A in SpA in the hope of furthering our understanding and improving treatment of these diseases.

Correction notice This article has been corrected since it published Online First. The last sentence in the second paragraph of the 'Well-defined role of IL-17A in host defence' section has been updated for clarity.

Acknowledgements The authors would like to acknowledge the following individuals for their help in the preparation of this work: Darren Asquith and Lindsey Hanson from Novartis UK.

Contributors All authors provided a substantial contribution to the conception, design and interpretation of the work, drafted the work or revised it critically for important intellectual content, and provided final approval of the submitted version of the manuscript.

Funding Medical writing support was provided by Ben Drever from Seren Communications, an Ashfield Company, part of UDG Healthcare, the funding for which was provided by Novartis.

Competing interests RM reports grants and personal fees from Novartis, during the conduct of the study. DGM reports grant and personal fees from Novartis, outside the submitted work. IBM reports personal fees from AbbVie, grants and personal fees from BMS, personal fees from Celgene, grants and personal fees from Janssen, personal fees from Lilly, personal fees from Novartis, grants and personal fees from UCB, personal fees from Leo, grants from Astra Zeneca, grants from Boehringer Ingelheim, outside the submitted work; BWK reports grants and personal fees from AbbVie, grants and personal fees from Eli Lilly & Co, grants and personal fees from Novartis, grants from Roche, grants from UCB, personal fees from Janssen, personal fees from Pfizer, personal fees from Sandoz, outside the submitted work. JS reports personal fees from AbbVie, personal fees from BMS, personal fees from Celgene, personal fees from Janssen, personal fees from Novartis, personal fees from UCB, personal fees from Merck, outside the submitted work.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Wright V. Seronegative polyarthritis: a unified concept. *Arthritis Rheum* 1978;21:619–33.
- Rudwaleit M, Khan MA, Sieper J. The challenge of diagnosis and classification in early ankylosing spondylitis: do we need new criteria? *Arthritis Rheum* 2005;52:1000–8.
- van der Heijde D, Ramiro S, Landewe R, et al. Update of the ASAS-EULAR management recommendations for axial spondyloarthritis. *Ann Rheum Dis* 2016;2017:978–91.
- McGonagle D, McDermott MF. A proposed classification of the immunological diseases. *PLoS Med* 2006;3:e297.
- Lande R, Botti E, Jandus C, et al. The antimicrobial peptide LL37 is a T-cell autoantigen in psoriasis. *Nat Commun* 2014;5.
- Mease PJ, Armstrong AW. Managing patients with psoriatic disease: the diagnosis and pharmacologic treatment of psoriatic arthritis in patients with psoriasis. *Drugs* 2014;74:423–41.
- Jadon DR, Sengupta R, Nightingale A, et al. Axial disease in psoriatic arthritis study: defining the clinical and radiographic phenotype of psoriatic spondyloarthritis. *Ann Rheum Dis* 2017;76:701–7.
- Brewerton DA, Nicholls A, Caffrey M, et al. HL-A 27 and arthropathies associated with ulcerative colitis and psoriasis. *The Lancet* 1974;303:956–8.
- Brown MA, Pile KD, Kennedy LG, et al. HLA class I associations of ankylosing spondylitis in the white population in the United Kingdom. *Ann Rheum Dis* 1996;55:268–70.
- Winchester R, Minevich G, Steshenko V, et al. HLA associations reveal genetic heterogeneity in psoriatic arthritis and in the psoriasis phenotype. *Arthritis Rheum* 2012;64:1134–44.
- Taams LS, Steel KJA, Srenathan U, et al. IL-17 in the immunopathogenesis of spondyloarthritis. *Nat Rev Rheumatol* 2018;14:453–66.
- Langley RG, Elewski BE, Lebwohl M, et al. Secukinumab in plaque psoriasis—results of two phase 3 trials. *N Engl J Med* 2014;371:326–38.
- Mease P, van der Heijde D, Landewe R, et al. Secukinumab improves active psoriatic arthritis symptoms and inhibits radiographic progression: primary results from the randomised, double-blind, phase III future 5 study. *Ann Rheum Dis* 2018;77:890–7.
- Mease PJ, McInnes IB, Kirkham B, et al. Secukinumab inhibition of interleukin-17A in patients with psoriatic arthritis. *N Engl J Med* 2015;373:1329–39.
- McInnes IB, Mease PJ, Kirkham B, et al. Secukinumab, a human anti-interleukin-17A monoclonal antibody, in patients with psoriatic arthritis (future 2): a randomised, double-blind, placebo-controlled, phase 3 trial. *The Lancet* 2015;386:1137–46.
- Baeten D, Sieper J, Braun J, et al. Secukinumab, an interleukin-17A inhibitor, in ankylosing spondylitis. *N Engl J Med* 2015;373:2534–48.
- Gordon KB, Blauvelt A, Papp KA, et al. Phase 3 trials of ixekizumab in moderate-to-severe plaque psoriasis. *N Engl J Med* 2016;375:345–56.
- Nash P, Kirkham B, Okada M, et al. Ixekizumab for the treatment of patients with active psoriatic arthritis and an inadequate response to tumour necrosis factor inhibitors: results from the 24-week randomised, double-blind, placebo-controlled period of the SPIRIT-P2 phase 3 trial. *The Lancet* 2017;389:2317–27.
- Mease PJ, van der Heijde D, Ritchlin CT, et al. Ixekizumab, an interleukin-17A specific monoclonal antibody, for the treatment of biologic-naïve patients with active psoriatic arthritis: results from the 24-week randomised, double-blind, placebo-controlled and active (adalimumab)-controlled period of the phase III trial SPIRIT-P1. *Ann Rheum Dis* 2017;76:79–87.
- Hueber W, Sands BE, Lewitzky S, et al. Secukinumab, a human anti-IL-17A monoclonal antibody, for moderate to severe Crohn's disease: unexpected results of a randomised, double-blind placebo-controlled trial. *Gut* 2012;61:1693–700.
- Dick AD, Tugal-Tutkun I, Foster S, et al. Secukinumab in the treatment of noninfectious uveitis: results of three randomized, controlled clinical trials. *Ophthalmology* 2013;120:777–87.
- Sharma SM, Fu DJ, Xue K. A review of the landscape of targeted immunomodulatory therapies for non-infectious uveitis. *Ophthalmol Ther* 2018;7:1–17.
- Gaffen SL. Structure and signalling in the IL-17 receptor family. *Nat Rev Immunol* 2009;9:556–67.
- Amatya N, Garg AV, Gaffen SL. IL-17 signaling: the yin and the Yang. *Trends Immunol* 2017;38:310–22.
- Ishigame H, Kakuta S, Nagai T, et al. Differential roles of interleukin-17A and -17F in host defense against mucocutaneous bacterial infection and allergic responses. *Immunity* 2009;30:108–19.
- Cho JS, Pietras EM, Garcia NC, et al. IL-17 is essential for host defense against cutaneous *Staphylococcus aureus* infection in mice. *J Clin Invest* 2010;120:1762–73.
- Aujla SJ, Dubin PJ, Kolls JK. Th17 cells and mucosal host defense. *Semin Immunol* 2007;19:377–82.
- Matsuzaki G, Umemura M. Interleukin-17 family cytokines in protective immunity against infections: role of hematopoietic cell-derived and non-hematopoietic cell-derived interleukin-17s. *Microbiology and Immunology* 2018;62:1–13.
- Lories RJ, Haroon N. Evolving concepts of new bone formation in axial spondyloarthritis: insights from animal models and human studies. *Best Practice & Research Clinical Rheumatology* 2017;31:877–86.
- Okada S, Puel A, Casanova J-L, et al. Chronic mucocutaneous candidiasis disease associated with inborn errors of IL-17 immunity. *Clin Transl Immunology* 2016;5:e114.
- Puel A, Cypowyj S, Maródi L, et al. Inborn errors of human IL-17 immunity underlie chronic mucocutaneous candidiasis. *Curr Opin Allergy Clin Immunol* 2012;12:616–22.

- 32 Mease PJ, McInnes IB, Reich K, *et al.* Secukinumab demonstrates a consistent safety profile with up to 5 years treatment in patients with psoriatic arthritis and moderate to severe plaque psoriasis: updated pooled safety analyses (Abstract THU0325). *Ann Rheum Dis* 2018;77:381–2.
- 33 Deodhar A, Baraliakos X, Marzo-Ortega H, *et al.* Secukinumab demonstrates consistent safety over long-term exposure (up to 3 years) in patients with active ankylosing spondylitis: pooled analysis of three phase 3 trials. *Arthritis Rheumatol* 2017;69.
- 34 Kammüller M, Tsai T-F, Griffiths CEM, *et al.* Inhibition of IL-17A by secukinumab shows no evidence of increased Mycobacterium tuberculosis infections. *Clin Transl Immunology* 2017;6:e152.
- 35 Deodhar A, Mease PJ, McInnes IB, *et al.* Long-term safety of secukinumab in patients with moderate-to-severe plaque psoriasis, psoriatic arthritis, and ankylosing spondylitis: integrated pooled clinical trial and post-marketing surveillance data. *Arthritis Res Ther* 2019;21.
- 36 Brown MA, Kenna T, Wordsworth BP. Genetics of ankylosing spondylitis—insights into pathogenesis. *Nat Rev Rheumatol* 2016;12:81–91.
- 37 Bowness P, Ridley A, Shaw J, *et al.* Th17 cells expressing KIR3DL2+ and responsive to HLA-B27 homodimers are increased in ankylosing spondylitis. *J Immunol* 2011;186:2672–80.
- 38 Allen RL, O'Callaghan CA, McMichael AJ, *et al.* Cutting edge: HLA-B27 can form a novel beta 2-microglobulin-free heavy chain homodimer structure. *J Immunol* 1999;162:5045–8.
- 39 DeLay ML, Turner MJ, Klenk EI, *et al.* HLA-B27 misfolding and the unfolded protein response augment interleukin-23 production and are associated with Th17 activation in transgenic rats. *Arthritis Rheum* 2009;60:2633–43.
- 40 Colbert RA, DeLay ML, Klenk EI, *et al.* From HLA-B27 to spondyloarthritis: a journey through the ER. *Immunol Rev* 2010;233:181–202.
- 41 Burton PR, Clayton DG, Cardon LR, *et al.* Association scan of 14,500 nonsynonymous SNPs in four diseases identifies autoimmunity variants. *Nat Genet* 2007;39:1329–37.
- 42 Evans DM, Spencer CCA, Pointon JJ, *et al.* Interaction between ERAP1 and HLA-B27 in ankylosing spondylitis implicates peptide handling in the mechanism for HLA-B27 in disease susceptibility. *Nat Genet* 2011;43:761–7.
- 43 Tsoi LC, Spain SL, Knight J, *et al.* Identification of 15 new psoriasis susceptibility loci highlights the role of innate immunity. *Nat Genet* 2012;44:1341–8.
- 44 Cortes A, Hadler J, Pointon JP, *et al.* Identification of multiple risk variants for ankylosing spondylitis through high-density genotyping of immune-related loci. *Nat Genet* 2013;45:730–8.
- 45 Filer C, Ho P, Smith RL, *et al.* Investigation of association of the IL12B and IL23R genes with psoriatic arthritis. *Arthritis Rheum* 2008;58:3705–9.
- 46 Bowes J, Orozco G, Flynn E, *et al.* Confirmation of TNIP1 and IL23a as susceptibility loci for psoriatic arthritis. *Ann Rheum Dis* 2011;70:1641–4.
- 47 Jostins L, Ripke S, Weersma RK, *et al.* Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature* 2012;491:119–24.
- 48 Coffre M, Roumier M, Rybczynska M, *et al.* Combinatorial control of Th17 and Th1 cell functions by genetic variations in genes associated with the interleukin-23 signaling pathway in spondyloarthritis. *Arthritis Rheum* 2013;65:1510–21.
- 49 de Beaucoudrey L, Puel A, Filipe-Santos O, *et al.* Mutations in STAT3 and IL12RB1 impair the development of human IL-17-producing T cells. *J Exp Med* 2008;205:1543–50.
- 50 Danoy P, Pryce K, Hadler J, *et al.* Association of variants at 1q32 and STAT3 with ankylosing spondylitis suggests genetic overlap with Crohn's disease. *PLoS Genet* 2010;6:e1001195.
- 51 Davidson SI, Liu Y, Danoy PA, *et al.* Association of STAT3 and TNFRSF1A with ankylosing spondylitis in Han Chinese. *Ann Rheum Dis* 2011;70:289–92.
- 52 Cénit MC, Ortego-Centeno N, Raya E, *et al.* Influence of the STAT3 genetic variants in the susceptibility to psoriatic arthritis and Behçet's disease. *Hum Immunol* 2013;74:230–3.
- 53 Harris TJ, Grosso JF, Yen H-R, *et al.* Cutting edge: an in vivo requirement for STAT3 signaling in Th17 development and TH17-dependent autoimmunity. *The Journal of Immunology* 2007;179:4313–7.
- 54 Soare A, Weber S, Maul L, *et al.* Cutting edge: homeostasis of innate lymphoid cells is imbalanced in psoriatic arthritis. *Ji* 2018;200:1249–54.
- 55 Leijten EFA, van Kempen TS, Boes M, *et al.* Brief report: enrichment of activated group 3 innate lymphoid cells in psoriatic arthritis synovial fluid. *Arthritis Rheumatol* 2015;67:2673–8.
- 56 Cuthbert R, Fragkakis E, Dunsmuir R, *et al.* FRI0427 first description of gamma delta T cells at normal human entheses. *Ann Rheum Dis* 2017;76:648.
- 57 Cuthbert R, Fragkakis EM, Bridgewood C, *et al.* The Vδ2 subset of TδT-cells are present at healthy human entheses and have transcriptional and functional characteristics consistent with a capacity for IL-17A production in response to IL-23. *Arthritis Rheumatol* 2018;70:1833–44.
- 58 Cuthbert RJ, Fragkakis EM, Dunsmuir R, *et al.* Brief report: group 3 innate lymphoid cells in human Enthesis. *Arthritis & Rheumatology* 2017;69:1816–22.
- 59 Schett G, Lories RJ, D'Agostino M-A, *et al.* Enthesitis: From pathophysiology to treatment. *Nat Rev Rheumatol* 2017;13:731–41.
- 60 Venken K, Jacques P, Mortier C, *et al.* RORγt inhibition selectively targets IL-17 producing iNKT and γδ-T cells enriched in spondyloarthritis patients. *Nat Commun* 2019;10.
- 61 Zhao M, Svensson MND, Venken K, *et al.* Altered thymic differentiation and modulation of arthritis by invariant NKT cells expressing mutant ZAP70. *Nat Commun* 2018;9.
- 62 Park CO, Kupper TS. The emerging role of resident memory T cells in protective immunity and inflammatory disease. *Nat Med* 2015;21:688–97.
- 63 Clark RA. Resident memory T cells in human health and disease. *Sci Transl Med* 2015;7:269rv1–1.
- 64 Cheuk S, Schlums H, Gallais Sérézal I, *et al.* CD49a Expression Defines Tissue-Resident CD8+ T Cells Poised for Cytotoxic Function in Human Skin. *Immunity* 2017;46:287–300.
- 65 Kirkham BM, Menon B, Taams LS. Immunological and clinical relationships of synovial IL-17+ T cells in psoriatic arthritis. *ACR/ARHP Annual Meeting* 2014;633.
- 66 Gaston JSH, Jadon DR. Th17 cell responses in spondyloarthritis. *Best Pract Res Clin Rheumatol* 2017;31:777–96.
- 67 Lynde CW, Poulin Y, Vender R, *et al.* Interleukin 17A: toward a new understanding of psoriasis pathogenesis. *J Am Acad Dermatol* 2014;71:141–50.
- 68 Coimbra S, Figueiredo A, Castro E, *et al.* The roles of cells and cytokines in the pathogenesis of psoriasis. *Int J Dermatol* 2012;51:389–98. quiz 95–8.
- 69 Coimbra S, Oliveira H, Reis F, *et al.* Interleukin (IL)-22, IL-17, IL-23, IL-8, vascular endothelial growth factor and tumour necrosis factor-α levels in patients with psoriasis before, during and after psoralen-ultraviolet A and narrowband ultraviolet B therapy. *Br J Dermatol* 2010;163:1282–90.
- 70 Lowes MA, Kikuchi T, Fuentes-Duculan J, *et al.* Psoriasis vulgaris lesions contain discrete populations of Th1 and Th17 T cells. *J Invest Dermatol* 2008;128:1207–11.
- 71 Menon B, Gullick NJ, Walter GJ, *et al.* Interleukin-17+CD8+ T cells are enriched in the joints of patients with psoriatic arthritis and correlate with disease activity and joint damage progression. *Arthritis Rheumatol* 2014;66:1272–81.
- 72 Jandus C, Bioley G, Rivals J-P, *et al.* Increased numbers of circulating polyfunctional Th17 memory cells in patients with seronegative spondylarthritides. *Arthritis Rheum* 2008;58:2307–17.
- 73 Shen H, Goodall JC, Hill Gaston JS. Frequency and phenotype of peripheral blood Th17 cells in ankylosing spondylitis and rheumatoid arthritis. *Arthritis Rheum* 2009;60:1647–56.
- 74 Shen H, Goodall JC, Gaston JSH. Frequency and phenotype of T helper 17 cells in peripheral blood and synovial fluid of patients with reactive arthritis. *J Rheumatol* 2010;37:2096–9.
- 75 Al-Mossawi MH, Chen L, Fang H, *et al.* Unique transcriptome signatures and GM-CSF expression in lymphocytes from patients with spondyloarthritis. *Nat Commun* 2017;8.
- 76 Benedetti G, Miossec P. Interleukin 17 contributes to the chronicity of inflammatory diseases such as rheumatoid arthritis. *Eur J Immunol* 2014;44:339–47.
- 77 Burkett PR, Meyer zu Horste G, Kuchroo VK. Pouring fuel on the fire: Th17 cells, the environment, and autoimmunity. *J Clin Invest* 2015;125:2211–9.
- 78 Lee YK, Mukasa R, Hatton RD, *et al.* Developmental plasticity of Th17 and Treg cells. *Curr Opin Immunol* 2009;21:274–80.
- 79 Wang C, Liao Q, Hu Y, *et al.* T lymphocyte subset imbalances in patients contribute to ankylosing spondylitis. *Exp Ther Med* 2015;9:250–6.
- 80 Teunissen MBM, Yeremenko NG, Baeten DLP, *et al.* The IL-17A-producing CD8+ T-cell population in psoriatic lesional skin comprises mucosa-associated invariant T cells and conventional T cells. *J Invest Dermatol* 2014;134:2898–907.
- 81 Gracey E, Qiayum Z, Almaghlouth I, *et al.* IL-7 primes IL-17 in mucosal-associated invariant T (MAIT) cells, which contribute to the Th17-axis in ankylosing spondylitis. *Ann Rheum Dis* 2016;75:2124–32.
- 82 Hayashi E, Chiba A, Tada K, *et al.* Involvement of mucosal-associated invariant T cells in ankylosing spondylitis. *J Rheumatol* 2016;43:1695–703.
- 83 Res PCM, Piskin G, de Boer OJ, *et al.* Overrepresentation of IL-17A and IL-22 producing CD8 T cells in lesional skin suggests their involvement in the pathogenesis of psoriasis. *PLoS One* 2010;5:e14108.
- 84 Kim J, Krueger JG. The immunopathogenesis of psoriasis. *Dermatol Clin* 2015;33:13–23.
- 85 Korn T, Bettelli E, Oukka M, *et al.* IL-17 and Th17 cells. *Annu Rev Immunol* 2009;27:485–517.
- 86 Keijsers RRM, Joosten I, van Erp PEJ, *et al.* Cellular sources of IL-17 in psoriasis: a paradigm shift? *Exp Dermatol* 2014;23:799–803.
- 87 Moran EM, Heydrich R, Ng CT, *et al.* IL-17A expression is localised to both mononuclear and polymorphonuclear synovial cell infiltrates. *PLoS One* 2011;6:e24048.
- 88 Appel H, Maier R, Wu P, *et al.* Analysis of IL-17(+) cells in facet joints of patients with spondyloarthritis suggests that the innate immune pathway might be of greater relevance than the Th17-mediated adaptive immune response. *Arthritis Res Ther* 2011;13.

- 89 Tamassia N, Arruda-Silva F, Calzetti F, et al. A Reappraisal on the Potential Ability of Human Neutrophils to Express and Produce IL-17 Family Members *In Vitro*: Failure to Reproducibly Detect It. *Front Immunol* 2018;9.
- 90 Noordenbos T, Yeremenko N, Gofita I, et al. Interleukin-17-positive mast cells contribute to synovial inflammation in spondylarthritis. *Arthritis Rheum* 2012;64:99–109.
- 91 Noordenbos T, Blijdorp I, Chen S, et al. Human mast cells capture, store, and release Bioactive, exogenous IL-17A. *J Leukoc Biol* 2016;100:453–62.
- 92 Chen S, Noordenbos T, Blijdorp I, et al. Histologic evidence that mast cells contribute to local tissue inflammation in peripheral spondylarthritis by regulating interleukin-17A content. *Rheumatology* 2019;58:617–27.
- 93 McGonagle D, Gibbon W, Emery P. Classification of inflammatory arthritis by enthesitis. *The Lancet* 1998;352:1137–40.
- 94 Cambré I, Gaublomme D, Schryvers N, et al. Running promotes chronicity of arthritis by local modulation of complement activators and impairing T regulatory feedback loops. *Ann Rheum Dis* 2019;78:787–95.
- 95 Reveille JD. Genetics of spondylarthritis—beyond the MHC. *Nat Rev Rheumatol* 2012;8:296–304.
- 96 McGonagle D, Tan AL. The enthesitis in psoriatic arthritis. *Clin Exp Rheumatol* 2015;33(5 Suppl 93):S36–9.
- 97 Xavier RJ, Podolsky DK. Unravelling the pathogenesis of inflammatory bowel disease. *Nature* 2007;448:427–34.
- 98 Sherlock JP, Joyce-Shaikh B, Turner SP, et al. IL-23 induces spondylarthropathy by acting on ROR- γ t+ CD3+CD4-CD8- enthesal resident T cells. *Nat Med* 2012;18:1069–76.
- 99 Reinhardt A, Yeva T, Worbs T, et al. Interleukin-23-Dependent $\gamma\delta$ T cells produce interleukin-17 and accumulate in the Enthesis, aortic valve, and ciliary body in mice. *Arthritis Rheumatol* 2016;68:2476–86.
- 100 Ono T, Okamoto K, Nakashima T, et al. IL-17-producing $\gamma\delta$ T cells enhance bone regeneration. *Nat Commun* 2016;7.
- 101 Benham H, Rehaume LM, Hasnain SZ, et al. Interleukin-23 mediates the intestinal response to microbial β -1,3-glucan and the development of spondylarthritis pathology in SKG mice. *Arthritis Rheumatol* 2014;66:1755–67.
- 102 Ruutu M, Thomas G, Steck R, et al. β -glucan triggers spondylarthritis and Crohn's disease-like ileitis in SKG mice. *Arthritis Rheum* 2012;64:2211–22.
- 103 Braem K, Carter S, Lories RJ. Spontaneous arthritis and ankylosis in male DBA/1 mice: further evidence for a role of behavioral factors in "stress-induced arthritis". *Biol Proced Online* 2012;14.
- 104 Ebihara S, Date F, Dong Y, et al. Interleukin-17 is a critical target for the treatment of ankylosing enthesitis and psoriasis-like dermatitis in mice. *Autoimmunity* 2015;48:259–66.
- 105 Rehaume LM, Mondot S, Aguirre de Cárcer D, et al. ZAP-70 genotype disrupts the relationship between microbiota and host, leading to spondylarthritis and ileitis in SKG mice. *Arthritis Rheumatol* 2014;66:2780–92.
- 106 Khmaladze I, Kelka T, Guerard S, et al. Mannan induces ROS-regulated, IL-17A-dependent psoriasis arthritis-like disease in mice. *Proceedings of the National Academy of Sciences* 2014;111:E3669–E3678.
- 107 Yamamoto M, Nakajima K, Takaishi M, et al. Psoriatic inflammation facilitates the onset of arthritis in a mouse model. *J Invest Dermatol* 2015;135:445–53.
- 108 Cambré I, Gaublomme D, Burssens A, et al. Mechanical strain determines the site-specific localization of inflammation and tissue damage in arthritis. *Nat Commun* 2018;9.
- 109 De Wilde K, Martens A, Lambrecht S, et al. A20 inhibition of STAT1 expression in myeloid cells: a novel endogenous regulatory mechanism preventing development of enthesitis. *Ann Rheum Dis* 2017;76:585–92.
- 110 Jacques P, Lambrecht S, Verheugen E, et al. Proof of concept: enthesitis and new bone formation in spondylarthritis are driven by mechanical strain and stromal cells. *Ann Rheum Dis* 2014;73:437–45.
- 111 Bridgewood C, Watad A, Russell T, et al. Identification of myeloid cells in the human enthesitis as the main source of local IL-23 production. *Ann Rheum Dis* 2019;78:929–33.
- 112 Yu JJ, Ruddy MJ, Wong GC, et al. An essential role for IL-17 in preventing pathogen-initiated bone destruction: recruitment of neutrophils to inflamed bone requires IL-17 receptor-dependent signals. *Blood* 2007;109:3794–802.
- 113 Ye P, Rodriguez FH, Kanaly S, et al. Requirement of interleukin 17 receptor signaling for lung CXCL chemokine and granulocyte colony-stimulating factor expression, neutrophil recruitment, and host defense. *J Exp Med* 2001;194:519–28.
- 114 Lories RJU, Luyten FP, de Vlam K. Progress in spondylarthritis. Mechanisms of new bone formation in spondylarthritis. *Arthritis Res Ther* 2009;11.
- 115 McGonagle D, Wakefield RJ, Tan AL, et al. Distinct topography of erosion and new bone formation in Achilles tendon enthesitis: implications for understanding the link between inflammation and bone formation in spondylarthritis. *Arthritis Rheum* 2008;58:2694–9.
- 116 van Duivenvoorde LM, Dorris ML, Satumira N, et al. Relationship between inflammation, bone destruction, and osteoproliferation in the HLA-B27/human β 2-microglobulin-transgenic rat model of spondylarthritis. *Arthritis Rheum* 2012;64:3210–9.
- 117 Dibra D, Xia X, Gagea M, et al. A spontaneous model of spondylarthropathies that develops bone loss and pathological bone formation: a process regulated by IL27RA-/- and mutant-p53. *PLoS One* 2018;13:e0193485.
- 118 Chabaud M, Lubberts E, Joosten L, et al. IL-17 derived from juxta-articular bone and synovium contributes to joint degradation in rheumatoid arthritis. *Arthritis Res* 2001;3:168–77.
- 119 Chabaud M, Miossec P. The combination of tumor necrosis factor γ blockade with interleukin-1 and interleukin-17 blockade is more effective for controlling synovial inflammation and bone resorption in an ex vivo model. *Arthritis & Rheumatism* 2001;44:1293–303.
- 120 Lubberts E, van den Bersselaar L, Oppers-Walgreen B, et al. IL-17 promotes bone erosion in murine collagen-induced arthritis through loss of the receptor activator of NF- κ B ligand/osteoprotegerin balance. *J Immunol* 2003;170:2655–62.
- 121 Adamopoulos IE, Chao C-chi, Geissler R, et al. Interleukin-17A upregulates receptor activator of NF- κ B on osteoclast precursors. *Arthritis Research & Therapy* 2010;12.
- 122 Sato K, Suematsu A, Okamoto K, et al. Th17 functions as an osteoclastogenic helper T cell subset that links T cell activation and bone destruction. *J Exp Med* 2006;203:2673–82.
- 123 Iwakura Y, Nakae S, Saijo S, et al. The roles of IL-17A in inflammatory immune responses and host defense against pathogens. *Immunol Rev* 2008;226:57–79.
- 124 Kotake S, Udagawa N, Takahashi N, et al. IL-17 in synovial fluids from patients with rheumatoid arthritis is a potent stimulator of osteoclastogenesis. *J. Clin. Invest.* 1999;103:1345–52.
- 125 Ke D, Fu X, Xue Y, et al. IL-17A regulates the autophagic activity of osteoclast precursors through RANKL-JNK1 signaling during osteoclastogenesis in vitro. *Biochem Biophys Res Commun* 2018;497:890–6.
- 126 Yang L, Fanok MH, Mediero-Munoz A, et al. Augmented Th17 differentiation leads to cutaneous and Synovio-Enthesal inflammation in a novel model of psoriatic arthritis. *Arthritis Rheumatol* 2018;70:855–67.
- 127 Wang Z, Jia Y, Du F, et al. IL-17A inhibits osteogenic differentiation of bone mesenchymal stem cells via Wnt signaling pathway. *Med Sci Monit* 2017;23:4095–101.
- 128 Zhang J-R, Pang D-D, Tong Q, et al. Different modulatory effects of IL-17, IL-22, and IL-23 on osteoblast differentiation. *Mediators of Inflammation* 2017;2017:1–11.
- 129 Mease PJ, Kavanaugh A, Reimold A, et al. Secukinumab in the treatment of psoriatic arthritis: efficacy and safety results through 3 years from the year 1 extension of the randomised phase III future 1 trial. *RMD Open* 2018;4:e000723.
- 130 Kampylafka E, d'Oliveira I, Linz C, et al. Resolution of synovitis and arrest of catabolic and anabolic bone changes in patients with psoriatic arthritis by IL-17A blockade with secukinumab: results from the prospective PSARTROS study. *Arthritis Res Ther* 2018;20.
- 131 Jo S, Wang SE, Lee YL, et al. IL-17A induces osteoblast differentiation by activating JAK2/STAT3 in ankylosing spondylitis. *Arthritis Res Ther* 2018;20.
- 132 Jo S, Lee JK, Han J, et al. Identification and characterization of human bone-derived cells. *Biochem Biophys Res Commun* 2018;495:1257–63.
- 133 Croes M, Öner FC, van Neerven D, et al. Proinflammatory T cells and IL-17 stimulate osteoblast differentiation. *Bone* 2016;84:262–70.
- 134 van Tok MN, van Duivenvoorde LM, Kramer I, et al. Interleukin-17A inhibition diminishes inflammation and new bone formation in experimental spondylarthritis. *Arthritis Rheumatol* 2019;71:612–25.
- 135 Ono T, Takayanagi H. Osteoimmunology in bone fracture healing. *Curr Osteoporosis Rep* 2017;15:367–75.
- 136 Uluçkan Özgü, Jimenez M, Karbach S, et al. Chronic skin inflammation leads to bone loss by IL-17-mediated inhibition of Wnt signaling in osteoblasts. *Sci Transl Med* 2016;8:330ra37–37.
- 137 Kim Y-G, Park J-W, Lee J-M, et al. IL-17 inhibits osteoblast differentiation and bone regeneration in rat. *Arch Oral Biol* 2014;59:897–905.
- 138 Ji R-R, Chamesian A, Zhang Y-Q. Pain regulation by non-neuronal cells and inflammation. *Science* 2016;354:572–7.
- 139 Pinho-Ribeiro FA, Verri WA, Chiu IM. Nociceptor sensory Neuron-Immune interactions in pain and inflammation. *Trends Immunol* 2017;38:5–19.
- 140 Blachier M, Canoui-Poitrine F, Dougados M, et al. Factors associated with radiographic lesions in early axial spondylarthritis. Results from the DESIR cohort. *Rheumatology* 2013;52:1686–93.
- 141 Wu Q, Inman RD, Davis KD. Neuropathic pain in ankylosing spondylitis: a psychophysics and brain imaging study. *Arthritis & Rheumatism* 2013;65:1494–503.
- 142 Ramjeeawon A, Choy E. 179 Neuropathic-like pain is common in psoriatic arthritis. *Rheumatology* 2018;57(suppl_3). key075.403-key075.403.
- 143 Richter F, Natta G, Ebbinghaus M, et al. Interleukin-17 sensitizes joint nociceptors to mechanical stimuli and contributes to arthritic pain through neuronal interleukin-17 receptors in rodents. *Arthritis Rheum* 2012;64:4125–34.
- 144 Pinto LG, Cunha TM, Vieira SM, et al. IL-17 mediates articular hypernociception in antigen-induced arthritis in mice. *Pain* 2010;148:247–56.
- 145 McNamee KE, Alzabin S, Hughes JP, et al. IL-17 induces hyperalgesia via TNF-dependent neutrophil infiltration. *Pain* 2011;152:1838–45.
- 146 Meng X, Zhang Y, Lao L, et al. Spinal interleukin-17 promotes thermal hyperalgesia and NMDA NR1 phosphorylation in an inflammatory pain rat model. *Pain* 2013;154:294–305.

- 147 Das Sarma J, Ciric B, Marek R, *et al.* Functional interleukin-17 receptor A is expressed in central nervous system glia and upregulated in experimental autoimmune encephalomyelitis. *J Neuroinflammation* 2009;6.
- 148 Hu Y, Ota N, Peng I, *et al.* IL-17RC is required for IL-17A- and IL-17F-dependent signaling and the pathogenesis of experimental autoimmune encephalomyelitis. *J Immunol* 2010;184:4307–16.
- 149 Cao L, DeLeo JA. CNS-infiltrating CD4+ T lymphocytes contribute to murine spinal nerve transection-induced neuropathic pain. *Eur J Immunol* 2008;38:448–58.
- 150 Costigan M, Moss A, Latremoliere A, *et al.* T-cell infiltration and signaling in the adult dorsal spinal cord is a major contributor to neuropathic pain-like hypersensitivity. *J Neurosci* 2009;29:14415–22.
- 151 Olechowski CJ, Truong JJ, Kerr BJ. Neuropathic pain behaviours in a chronic-relapsing model of experimental autoimmune encephalomyelitis (EAE). *Pain* 2009;141:156–64.
- 152 Sun C, Zhang J, Chen L, *et al.* IL-17 contributed to the neuropathic pain following peripheral nerve injury by promoting astrocyte proliferation and secretion of proinflammatory cytokines. *Mol Med Rep* 2017;15:89–96.
- 153 Deodhar A, Conaghan PG, Kvien TK, *et al.* Secukinumab provides rapid and persistent relief in pain and fatigue symptoms in patients with ankylosing spondylitis irrespective of baseline C-reactive protein levels or prior tumour necrosis factor inhibitor therapy: 2-year data from the measure 2 study. *Clin Exp Rheumatol* 2019;37:260–9.
- 154 McInnes IB, Mease PJ, Schett G, *et al.* Secukinumab provides rapid and sustained pain relief in psoriatic arthritis over 2 years: results from the FUTURE 2 study. *Arthritis Res Ther* 2018;20.
- 155 Hunderforan G, Neurath MF, Mudter J. Functional relevance of T helper 17 (Th17) cells and the IL-17 cytokine family in inflammatory bowel disease. *Inflamm Bowel Dis* 2012;18:180–6.
- 156 Khanna PV, Shih DQ, Haritunians T, *et al.* Use of animal models in elucidating disease pathogenesis in IBD. *Semin Immunopathol* 2014;36:541–51.
- 157 Schreiber S, Colombel J-F, Feagan BG, *et al.* Incidence rates of inflammatory bowel disease in patients with psoriasis, psoriatic arthritis and ankylosing spondylitis treated with secukinumab: a retrospective analysis of pooled data from 21 clinical trials. *Ann Rheum Dis* 2019;78:473–9.
- 158 Lee JS, Tato CM, Joyce-Shaikh B, *et al.* Interleukin-23-independent IL-17 production regulates intestinal epithelial permeability. *Immunity* 2015;43:727–38.
- 159 Ivanov II, Atarashi K, Manel N, *et al.* Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell* 2009;139:485–98.
- 160 Niess JH, Leithäuser F, Adler G, *et al.* Commensal gut flora drives the expansion of proinflammatory CD4 T cells in the colonic lamina propria under normal and inflammatory conditions. *J Immunol* 2008;180:559–68.
- 161 Mortier C, Govindarajan S, Venken K, *et al.* It Takes "Guts" to Cause Joint Inflammation: Role of Innate-Like T Cells. *Front Immunol* 2018;9.
- 162 Shih VF-S, Cox J, Kljavin NM, *et al.* Homeostatic IL-23 receptor signaling limits Th17 response through IL-22-mediated containment of commensal microbiota. *Proceedings of the National Academy of Sciences* 2014;111:13942–7.
- 163 Qiu J, Guo X, Chen Z-ME, *et al.* Group 3 innate lymphoid cells inhibit T-cell-mediated intestinal inflammation through aryl hydrocarbon receptor signaling and regulation of microflora. *Immunity* 2013;39:386–99.
- 164 Ciccio F, Guggino G, Rizzo A, *et al.* Type 3 innate lymphoid cells producing IL-17 and IL-22 are expanded in the gut, in the peripheral blood, synovial fluid and bone marrow of patients with ankylosing spondylitis. *Ann Rheum Dis* 2015;74:1739–47.
- 165 Manfredo Vieira S, Hiltensperger M, Kumar V, *et al.* Translocation of a gut pathobiont drives autoimmunity in mice and humans. *Science* 2018;359:1156–61.
- 166 Weinstein JE, Peppel KL. Cytokines in uveitis. *Curr Opin Ophthalmol* 2018;29:267–74.
- 167 Deodhar A, Miceli-Richard C, Baraliakos X, *et al.* Low incidence of both new-onset and flares of uveitis in secukinumab-treated patients with ankylosing spondylitis: clinical trial and post-marketing safety analysis (Abstract SAT0270). *Ann Rheum Dis* 2018;77:999.
- 168 Taçi D, Blauvelt A, Reich K, *et al.* Secukinumab is superior to ustekinumab in clearing skin of subjects with moderate to severe plaque psoriasis: clear, a randomized controlled trial. *J Am Acad Dermatol* 2015;73:400–9.
- 169 van der Heijde D, Cheng-Chung Wei J, Dougados M, *et al.* Ixekizumab, an interleukin-17A antagonist in the treatment of ankylosing spondylitis or radiographic axial spondyloarthritis in patients previously untreated with biological disease-modifying anti-rheumatic drugs (COAST-V): 16 week results of a phase 3 randomised, double-blind, active-controlled and placebo-controlled trial. *The Lancet* 2018;392:2441–51.
- 170 Deodhar A, Poddubnyy D, Pacheco-Tena C, *et al.* Efficacy and Safety of Ixekizumab in the Treatment of Radiographic Axial Spondyloarthritis: Sixteen-Week Results From a Phase III Randomized, Double-Blind, Placebo-Controlled Trial in Patients With Prior Inadequate Response to or Intolerance of Tumor Necrosis Factor Inhibitors. *Arthritis Rheumatol* 2019;71:599–611.
- 171 Deodhar A, Gensler LS, Sieper J, *et al.* Three multicenter, randomized, double-blind, placebo-controlled studies evaluating the efficacy and safety of ustekinumab in axial spondyloarthritis. *Arthritis Rheumatol* 2019;71:258–70.
- 172 Baeten D, Østergaard M, Wei JC-C, *et al.* Risankizumab, an IL-23 inhibitor, for ankylosing spondylitis: results of a randomised, double-blind, placebo-controlled, proof-of-concept, dose-finding phase 2 study. *Ann Rheum Dis* 2018;77:1295–302.
- 173 McGonagle D, Lories RJU, Tan AL, *et al.* The concept of a "synovio-entheseal complex" and its implications for understanding joint inflammation and damage in psoriatic arthritis and beyond. *Arthritis Rheum* 2007;56:2482–91.
- 174 Castillo-Gallego C, Aydin SZ, Emery P, *et al.* Magnetic resonance imaging assessment of axial psoriatic arthritis: extent of disease relates to HLA-B27. *Arthritis Rheum* 2013;65:2274–8.
- 175 van Tok MN, Na S, Lao CR, *et al.* The initiation, but not the persistence, of experimental spondyloarthritis is dependent on interleukin-23 signaling. *Front Immunol* 2018;9.
- 176 Fieschi C, Bosticardo M, de Beaucoudrey L, *et al.* A novel form of complete IL-12/IL-23 receptor beta1 deficiency with cell surface-expressed nonfunctional receptors. *Blood* 2004;104:2095–101.
- 177 Fieschi C, Casanova J-L. The role of interleukin-12 in human infectious diseases: only a faint signature. *Eur J Immunol* 2003;33:1461–4.
- 178 Fieschi C, Dupuis S, Catherinot E, *et al.* Low penetrance, broad resistance, and favorable outcome of interleukin 12 receptor beta1 deficiency: medical and immunological implications. *J Exp Med* 2003;197:527–35.
- 179 Ouederni M, Sanal O, Ikinciogullari A, *et al.* Clinical features of candidiasis in patients with inherited interleukin 12 receptor β 1 deficiency. *Clin Infect Dis* 2014;58:204–13.
- 180 Ramirez-Alejo N, Blancas-Galicia L, Yamazaki-Nakashimada M, *et al.* Molecular analysis for patients with IL-12 receptor β 1 deficiency. *Clin Genet* 2014;86:161–6.
- 181 Bustamante J, Boisson-Dupuis S, Jouanguy E, *et al.* Novel primary immunodeficiencies revealed by the investigation of paediatric infectious diseases. *Curr Opin Immunol* 2008;20:39–48.
- 182 Boisson B, Wang C, Pedergrana V, *et al.* An ACT1 mutation selectively abolishes interleukin-17 responses in humans with chronic mucocutaneous candidiasis. *Immunity* 2013;39:676–86.
- 183 Boisson-Dupuis S, Kong X-F, Okada S, *et al.* Inborn errors of human STAT1: Allelic heterogeneity governs the diversity of immunological and infectious phenotypes. *Curr Opin Immunol* 2012;24:364–78.
- 184 Altare F, Durandy A, Lammass D, *et al.* Impairment of mycobacterial immunity in human interleukin-12 receptor deficiency. *Science* 1998;280:1432–5.
- 185 Akahoshi M, Nakashima H, Miyake K, *et al.* Influence of interleukin-12 receptor beta1 polymorphisms on tuberculosis. *Hum Genet* 2003;112:237–43.
- 186 Al-Muhsen S, Casanova J-L. The genetic heterogeneity of Mendelian susceptibility to mycobacterial diseases. *J Allergy Clin Immunol* 2008;122:1043–51.
- 187 de Beaucoudrey L, Samarina A, Bustamante J, *et al.* Revisiting human IL-12R β 1 deficiency: a survey of 141 patients from 30 countries. *Medicine* 2010;89:381–402.
- 188 MacLennan C, Fieschi C, Lammass DA, *et al.* Interleukin (IL)-12 and IL-23 are key cytokines for immunity against Salmonella in humans. *J Infect Dis* 2004;190:1755–7.
- 189 Godínez I, Keestra AM, Spees A, *et al.* The IL-23 axis in Salmonella gastroenteritis. *Cell Microbiol* 2011;13:1639–47.
- 190 de Jong R, Altare F, Haagen IA, *et al.* Severe mycobacterial and Salmonella infections in interleukin-12 receptor-deficient patients. *Science* 1998;280:1435–8.
- 191 Puel A, Cypowyj S, Bustamante J, *et al.* Chronic mucocutaneous candidiasis in humans with inborn errors of interleukin-17 immunity. *Science* 2011;332:65–8.
- 192 Ling Y, Cypowyj S, Aytakin C, *et al.* Inherited IL-17RC deficiency in patients with chronic mucocutaneous candidiasis. *J Exp Med* 2015;212:619–31.
- 193 Bader O, Weig MS, Gross U, *et al.* Photo quiz. A 32-year-old man with ulcerative mucositis, skin lesions, and nail dystrophy. Chronic mucocutaneous candidiasis by multidrug-resistant *Candida albicans*. *Clin Infect Dis* 2012;54:1035–6.
- 194 Kisan K, Bøe Wolff AS, Podkrajšek KT, *et al.* Chronic mucocutaneous candidiasis in APECED or thymoma patients correlates with autoimmunity to Th17-associated cytokines. *J Exp Med* 2010;207:299–308.
- 195 Kisan K, Lilic D, Casanova J-L, *et al.* Mucocutaneous candidiasis and autoimmunity against cytokines in APECED and thymoma patients: clinical and pathogenetic implications. *Eur J Immunol* 2011;41:1517–27.
- 196 Aaltonen J, Björnsen P, Perheentupa J, *et al.* An autoimmune disease, APECED, caused by mutations in a novel gene featuring two PHD-type zinc-finger domains. *Nat Genet* 1997;17:399–403.
- 197 Nagamine K, Peterson P, Scott HS, *et al.* Positional cloning of the APECED gene. *Nat Genet* 1997;17:393–8.
- 198 Freeman AF, Holland SM. The hyper-IgE syndromes. *Immunol Allergy Clin North Am* 2008;28:727–91.
- 199 Renner ED, Rylaarsdam S, Añover-Sombke S, *et al.* Novel signal transducer and activator of transcription 3 (STAT3) mutations, reduced T(H)17 cell numbers, and variably defective STAT3 phosphorylation in hyper-IgE syndrome. *J Allergy Clin Immunol* 2008;122:181–7.
- 200 Ma CS, Chew GYJ, Simpson N, *et al.* Deficiency of Th17 cells in hyper IgE syndrome due to mutations in STAT3. *J Exp Med* 2008;205:1551–7.
- 201 Minegishi Y, Saito M, Tsuchiya S, *et al.* Dominant-negative mutations in the DNA-binding domain of STAT3 cause hyper-IgE syndrome. *Nature* 2007;448:1058–62.

- 202 Drewniak A, Gazendam RP, Tool ATJ, *et al.* Invasive fungal infection and impaired neutrophil killing in human CARD9 deficiency. *Blood* 2013;121:2385–92.
- 203 Engelhardt KR, McGhee S, Winkler S, *et al.* Large deletions and point mutations involving the dedicator of cytokinesis 8 (DOCK8) in the autosomal-recessive form of hyper-IgE syndrome. *J Allergy Clin Immunol* 2009;124:1289–302.
- 204 Lee YH, Song GG. Associations between interleukin-23R polymorphisms and ankylosing spondylitis susceptibility: an updated meta-analysis. *Z Rheumatol* 2019;78:272–80.
- 205 Yang B, Xu Y, Liu X, *et al.* IL-23R and IL-17A polymorphisms correlate with susceptibility of ankylosing spondylitis in a southwest Chinese population. *Oncotarget* 2017;8:70310–6.
- 206 Zhong L, Wang W, Song H. Complex role of IL-23R polymorphisms on ankylosing spondylitis: a meta-analysis. *Expert Rev Clin Immunol* 2018;14:635–43.
- 207 Chen C, Zhang X, Wang Y. ANTXR2 and IL-1R2 polymorphisms are not associated with ankylosing spondylitis in Chinese Han population. *Rheumatol Int* 2012;32:15–19.
- 208 Rocha Loures MA, Macedo LC, Reis DM, *et al.* Influence of *TNF* and *IL17* Gene Polymorphisms on the Spondyloarthritis Immunopathogenesis, Regardless of HLA-B27, in a Brazilian Population. *Mediators Inflamm* 2018;2018:1–7.
- 209 Vidal-Castieira JR, López-Vázquez A, Díaz-Peña R, *et al.* A single nucleotide polymorphism in the *IL17ra* promoter is associated with functional severity of ankylosing spondylitis. *PLoS One* 2016;11:e0158905.
- 210 Nossent JC, Sagen-Johnsen S, Bakland G. IL23R gene variants in relation to IL17A levels and clinical phenotype in patients with ankylosing spondylitis. *Rheumatol Adv Pract* 2018;2.
- 211 Reveille JD, Sims A-M, Danoy P, *et al.* Genome-wide association study of ankylosing spondylitis identifies non-MHC susceptibility loci. *Nat Genet* 2010;42:123–7.
- 212 Karaderi T, Harvey D, Farrar C, *et al.* Association between the interleukin 23 receptor and ankylosing spondylitis is confirmed by a new UK case-control study and meta-analysis of published series. *Rheumatology* 2009;48:386–9.
- 213 Dong H, Li Q, Zhang Y, *et al.* IL23R gene confers susceptibility to ankylosing spondylitis concomitant with uveitis in a Han Chinese population. *PLoS One* 2013;8:e67505.
- 214 Wang X, Huang J, Lin Z, *et al.* Single-nucleotide polymorphisms and expression of IL23R in Chinese ankylosing spondylitis patients. *Rheumatol Int* 2010;30:955–9.
- 215 Sáfrány E, Pászár B, Csöngéi V, *et al.* Variants of the IL23R gene are associated with ankylosing spondylitis but not with Sjögren syndrome in Hungarian population samples. *Scand J Immunol* 2009;70:68–74.
- 216 Roberts AR, Vecellio M, Cortes A, *et al.* Investigation of a possible extended risk haplotype in the IL23R region associated with ankylosing spondylitis. *Genes Immun* 2017;18:105–8.
- 217 Xia Y, Liu Y-Q, Chen K, *et al.* Association of IL-1R2 genetic polymorphisms with the susceptibility of ankylosing spondylitis in northern Chinese Han population. *Mod Rheumatol* 2015;25:908–12.
- 218 Kenna TJ, Davidson SJ, Duan R, *et al.* Enrichment of circulating interleukin-17-secreting interleukin-23 receptor-positive $\gamma\delta$ T cells in patients with active ankylosing spondylitis. *Arthritis Rheum* 2012;64:1420–9.
- 219 Guggino G, Ciccía F, Di Liberto D, *et al.* Interleukin (IL)-9/IL-9R axis drives $\gamma\delta$ T cells activation in psoriatic arthritis patients. *Clin Exp Immunol* 2016;186:277–83.
- 220 Chowdhury AC, Chaurasia S, Mishra SK, *et al.* IL-17 and IFN- γ producing NK and $\gamma\delta$ -T cells are preferentially expanded in synovial fluid of patients with reactive arthritis and undifferentiated spondyloarthritis. *Clin Immunol* 2017;183:207–12.
- 221 Cai Y, Shen X, Ding C, *et al.* Pivotal role of dermal IL-17-producing $\gamma\delta$ T cells in skin inflammation. *Immunity* 2011;35:596–610.
- 222 Campbell JJ, Ebsworth K, Ertl LS, *et al.* IL-17–Secreting $\gamma\delta$ T cells are completely dependent upon CCR6 for homing to inflamed skin. *J. i.* 2017;199:3129–36.
- 223 Gravallese EM, Schett G. Effects of the IL-23-IL-17 pathway on bone in spondyloarthritis. *Nat Rev Rheumatol* 2018;14:631–40.
- 224 Leipe J, Grunke M, Dechant C, *et al.* Role of Th17 cells in human autoimmune arthritis. *Arthritis Rheum* 2010;62:2876–85.
- 225 Xueyi L, Lina C, Zhenbiao W, *et al.* Levels of circulating Th17 cells and regulatory T cells in ankylosing spondylitis patients with an inadequate response to anti-TNF- α therapy. *J Clin Immunol* 2013;33:151–61.
- 226 Zizzo G, De Santis M, Bosello SL, *et al.* Synovial fluid-derived T helper 17 cells correlate with inflammatory activity in arthritis, irrespectively of diagnosis. *Clin Immunol* 2011;138:107–16.
- 227 Ortega C, Fernández-A S, Carrillo JM, *et al.* IL-17-producing CD8⁺ T lymphocytes from psoriasis skin plaques are cytotoxic effector cells that secrete Th17-related cytokines. *J Leukoc Biol* 2009;86:435–43.
- 228 Hijnen D, Knol EF, Gent YY, *et al.* CD8(+) T cells in the lesional skin of atopic dermatitis and psoriasis patients are an important source of IFN- γ , IL-13, IL-17, and IL-22. *J Invest Dermatol* 2013;133:973–9.
- 229 Steel KJA, S-Y W, Srenathan U, *et al.* Synovial IL-17+ CD8+ T cells are a pro-inflammatory tissue resident population enriched in spondyloarthritis (Abstract 0016). *Ann Rheum Dis* 2018;77:A8–A9.
- 230 Di Meglio P, Villanova F, Navarini AA, *et al.* Targeting CD8(+) T cells prevents psoriasis development. *J Allergy Clin Immunol* 2016;138:274–6.
- 231 Rahman MS, Akhtar N, Jamil HM, *et al.* TGF- β /BMP signaling and other molecular events: regulation of osteoblastogenesis and bone formation. *Bone Res* 2015;3.
- 232 ClinicalTrialsgov. *Study of the efficacy and safety of Secukinumab in participants with active psoriatic arthritis with axial skeleton involvement (maximise)*. National Institute of Health US National Library of Medicine, 2018.
- 233 Braun J, Baraliakos X, Deodhar A, *et al.* Effect of secukinumab on clinical and radiographic outcomes in ankylosing spondylitis: 2-year results from the randomised phase III measure 1 study. *Ann Rheum Dis* 2017;76:1070–7.
- 234 ClinicalTrialsgov. *NCT02696031: study of efficacy and safety of Secukinumab in patients with non-radiographic axial spondyloarthritis*. National Institute of Health US National Library of Medicine, 2018.
- 235 Novartis. Cosentyx: EPAR - Product Information, 2017. Available: http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/003729/human_med_001832.jsp&mid=WC0b01ac058001d124
- 236 ClinicalTrialsgov. *NCT02757352: a study of ixekizumab (LY2439821) in participants with Nonradiographic axial spondyloarthritis (COAST-X)*, 2018. National Institute of health us National Library of medicine. Available: <https://clinicaltrials.gov/ct2/show/NCT02757352>
- 237 Lilly. Taltz EPAR, 2019. Available: https://www.ema.europa.eu/documents/product-information/taltz-epar-product-information_en.pdf
- 238 Sieper J, Braun J, Dougados M, *et al.* Axial spondyloarthritis. *Nat Rev Dis Primers* 2015;1.
- 239 McInnes IB, Kavanaugh A, Gottlieb AB, *et al.* Efficacy and safety of ustekinumab in patients with active psoriatic arthritis: 1 year results of the phase 3, multicentre, double-blind, placebo-controlled PSUMMIT 1 trial. *Lancet* 2013;382:780–9.
- 240 Ritchlin C, Rahman P, Kavanaugh A, *et al.* Efficacy and safety of the anti-IL-12/23 p40 monoclonal antibody, ustekinumab, in patients with active psoriatic arthritis despite conventional non-biological and biological anti-tumour necrosis factor therapy: 6-month and 1-year results of the phase 3, multicentre, double-blind, placebo-controlled, randomised PSUMMIT 2 trial. *Ann Rheum Dis* 2014;73:990–9.
- 241 Leonardi CL, Kimball AB, Papp KA, *et al.* Efficacy and safety of ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with psoriasis: 76-week results from a randomised, double-blind, placebo-controlled trial (Phoenix 1). *The Lancet* 2008;371:1665–74.
- 242 Griffiths CEM, Strober BE, van de Kerkhof P, *et al.* Comparison of ustekinumab and etanercept for moderate-to-severe psoriasis. *N Engl J Med* 2010;362:118–28.
- 243 Reich K, Armstrong AW, Foley P, *et al.* Efficacy and safety of guselkumab, an anti-interleukin-23 monoclonal antibody, compared with adalimumab for the treatment of patients with moderate to severe psoriasis with randomized withdrawal and retreatment: results from the phase III, double-blind, placebo- and active comparator-controlled voyage 2 trial. *J Am Acad Dermatol* 2017;76:418–31.
- 244 Gordon KB, Strober B, Lebwohl M, *et al.* Efficacy and safety of risankizumab in moderate-to-severe plaque psoriasis (UltIMMA-1 and UltIMMA-2): results from two double-blind, randomised, placebo-controlled and ustekinumab-controlled phase 3 trials. *The Lancet* 2018;392:650–61.
- 245 Blauvelt A, Papp KA, Griffiths CEM, *et al.* Efficacy and safety of guselkumab, an anti-interleukin-23 monoclonal antibody, compared with adalimumab for the continuous treatment of patients with moderate to severe psoriasis: results from the phase III, double-blind, placebo- and active comparator-controlled voyage 1 trial. *J Am Acad Dermatol* 2017;76:405–17.
- 246 van der Heijde D, Kivitz A, Schiff MH, *et al.* Efficacy and safety of adalimumab in patients with ankylosing spondylitis: results of a multicenter, randomized, double-blind, placebo-controlled trial. *Arthritis Rheum* 2006;54:2136–46.
- 247 Mease PJ, Gladman DD, Ritchlin CT, *et al.* Adalimumab for the treatment of patients with moderately to severely active psoriatic arthritis: results of a double-blind, randomized, placebo-controlled trial. *Arthritis Rheum* 2005;52:3279–89.
- 248 Menter A, Tyring SK, Gordon K, *et al.* Adalimumab therapy for moderate to severe psoriasis: a randomized, controlled phase III trial. *J Am Acad Dermatol* 2008;58:106–15.
- 249 Yoshiga Y, Goto D, Segawa S, *et al.* Invariant NKT cells produce IL-17 through IL-23-dependent and -independent pathways with potential modulation of Th17 response in collagen-induced arthritis. *Int J Mol Med* 2008;22:369–74.